



Team  
Ophthalmologist



Team  
Ophthalmologist



**SCHWARTZ**  
LASER EYE CENTER



Team  
Lasik Surgeon

## CATARACT SURGERY / LENS EXCHANGE EYE MEDICATIONS

We will give you a written prescription for all of the following eye drops at your A-Scan (measurement) appointment. Please have filled at your pharmacy immediately after receiving your written prescriptions.



### 1. AZASITE (Antibiotic/Anti-inflammatory): **BEFORE SURGERY ONLY DROP**

- You will start using AzaSite 2 weeks prior to your surgery date. You will continue using AzaSite up until the night before surgery in your procedure eye.
- Take AzaSite 1-drop at night in procedure eye.

**DO NOT TAKE AZASITE THE DAY OF YOUR SURGERY. AZASITE WILL BE USED PRIOR TO PROCEDURE ONLY.**



### 2. ZYMAXID / Generic: GATIFLOXACIN (Antibiotic): **BEFORE & AFTER drop**

- You will start Zymaxid/Gatifloxacin 3 days BEFORE your surgery. Continue using this drop on the day of your procedure.
- Zymaxid / Gatifloxacin will be used 4 X per day in your procedure eye.
- Zymaxid / Gatifloxacin should be used 1-drop (4) times per day for 1 week AFTER your procedure and then discontinue but do not discard.



### 3. BROMSITE (Anti-inflammatory): **BEFORE & AFTER drop**

- You will start BromSite 3 days BEFORE your surgery including surgery. Continue using this drop on the day of your procedure.
- BromSite will be used 2 X per day in your procedure eye.
- BromSite should be used 1-drop (2) times per day for 4 weeks AFTER your procedure and then discontinue but do not discard.



### 4. PRED FORTE / Generic: Prednisolone (Steriod): **AFTER SX drop**

- Pred Forte / Prednisolone should be used 1-drop (4) times per day for one week, then 1-drop (3) times per day for one week, then 1-drop (2) times per day for one week, then 1-drop (1) times per day for one week. After completion of the four week tapering schedule discontinue but do not discard.

**SHAKE WELL BEFORE EACH USE**

• If your eye is **NOT** patched after surgery; Please restart your **BEFORE & AFTER (Zymaxid/Gatifloxacin, Bromsite, Pred Fote/ Prednisolone)** drops (3) hours AFTER your surgery.

• If your eye **IS PATCHED** after surgery; Leave your eye patched until the morning after your surgery. You can remove the patch and start your **BEFORE & AFTER (Zymaxid/Gatifloxacin, Bromsite, Pred Fote/ Prednisolone).**