



Team
Ophthalmologist



Team
Ophthalmologist



SCHWARTZ LASER EYE CENTER



Team
Lasik Surgeon

CATARACT SURGERY / LENS EXCHANGE BEFORE SURGERY MEDICATIONS

Start the medications listed below before your procedure in the OPERATIVE EYE.



HypoChlor Solution (Preventative Antibiotic): **BEFORE SURGERY ONLY** Cleanser

- You will start cleansing lids with HypoChlor 2 weeks prior to your surgery date. You will continue using HypoChlor up until the night before surgery on both **EYELIDS**.
- HypoChlor is a spray to be used with a cleaning cloth, cotton ball, ect.

Start using HypoChlor on _____, on Both EYELIDS

Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>
Day 8	Day 9	Day 10	Day 11	Day 12	Day 13	Day 14
<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>



ZYMAXID / Generic: GATIFLOXACIN (Antibiotic): **BEFORE & AFTER drop do not discard**

- You will start Zymaxid/Gatifloxacin 3 days BEFORE your surgery. Continue using this drop on the day of your procedure.
- Zymaxid / Gatifloxacin will be used 4 X per day in your procedure eye.
- Zymaxid / Gatifloxacin should be used 1-drop (4) times per day for 1 week AFTER your procedure and then discontinue but do not discard.

Start using Zymaxid/Gatifloxacin on _____, in your _____ EYE

3 DAYS PRIOR	2 DAYS PRIOR	1 DAY PRIOR	DAY OF PROCEDURE
<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>

Helpful Hint: Morning, Lunch, Dinner, Bedtime.

wait 5 minutes between drops * wait 5 minutes between drops *** wait 5 wait minutes between drops***



3. BROMSITE (Anti-inflammatory): **BEFORE & AFTER drop do not discard**

- You will start BromSite 3 days BEFORE your surgery including surgery. Continue using this drop on the day of your procedure.
- BromSite will be used 2 X per day in your procedure eye.
- BromSite should be used 1-drop (2) times per day for 4 weeks AFTER your procedure and then discontinue but do not discard.

Start using BromSite on _____, in your _____ EYE

3 DAYS PRIOR	2 DAYS PRIOR	1 DAY PRIOR	DAY OF PROCEDURE
<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/>

Helpful Hint: Morning & Bedtime.